

Pretty Good State Transfer in Qubit Chains - The Heisenberg Hamiltonian

Leonardo Banchi,¹ Gabriel Coutinho,^{2, a)} Chris Godsil,³ and Simone Severini⁴

¹⁾*Department of Physics and Astronomy, University College London,
London, WC1E 6BT, United Kingdom*

²⁾*Departamento de Ciência da Computação, IME-USP,
São Paulo, SP, 05508-090, Brazil*

³⁾*Department of Combinatorics & Optimization, University of Waterloo,
Waterloo, ON, N2L3G1, Canada*

⁴⁾*Department of Computer Science, University College London,
London, WC1E 6BT, United Kingdom*

(Dated: 23 February 2017)

Pretty good state transfer in networks of qubits occurs when a continuous-time quantum walk allows the transmission of a qubit state from one node of the network to another, with fidelity arbitrarily close to 1. We prove that in a Heisenberg chain with n qubits there is pretty good state transfer between the nodes at the j -th and $(n - j + 1)$ -th position if n is a power of 2. Moreover, this condition is also necessary for $j = 1$. We obtain this result by applying a theorem due to Kronecker about Diophantine approximations, together with techniques from algebraic graph theory.

^{a)}Electronic mail: coutinho@ime.usp.br

I. INTRODUCTION

Long-distance quantum communication, for example over several kilometers, typically uses photonic systems. On the other hand, given the difficulty of engineering interactions between photons, several promising candidates for quantum hardware are based on *quantum networks* of localized qubits¹, which are easier to manipulate. In typical quantum algorithms during the computation the states of different qubits have to be transferred between various registers, namely between different nodes of the network. However, when qubits are localized, their physical movement may be either impossible, by construction, or energy inefficient. A viable solution to this problem is to exploit the coherent dynamics of the quantum network, namely a continuous-time quantum walk, to transfer the quantum states between different nodes². This approach for short-distance (*in-chip*) communication has attracted much attention³ because it minimizes the use of external control and also avoids the complex interface between localized and moving particles. However, in a generic quantum network the resulting coherent dynamics is very complicated and the transmission between two nodes is normally inefficient. Therefore, much effort has been devoted to understand what are the best strategies, or the best networks, to achieve either *perfect state transfer* between distant nodes⁴⁻⁷, or *pretty good state transfer*⁸⁻¹⁶ where the transmission quality is *almost* perfect. One-dimensional systems, namely chains of qubits, are perhaps the most natural candidate for transmission as they resemble a quantum wire or *data-bus*.

Most of the literature on perfect or pretty good state transfer has considered chains interacting with the XY Hamiltonian (also called XX Hamiltonian in the condensed matter community¹⁷). This kind of chains offers several mathematical simplifications, e.g. the Hamiltonian in the single-particle subspace is equivalent to the adjacency matrix of the corresponding graph and also the many-particle problem is exactly solvable¹⁸. However, in solid state systems such as quantum dots¹⁹ or dopants in silicon²⁰ and in current optical lattice experiments²¹, which are some of the most promising quantum devices, the natural interaction is the Heisenberg (XYZ) Hamiltonian, sometimes also called XXX Hamiltonian¹⁷. For engineered chains in the single-particle subspace the distinction between XY and XYZ Hamiltonians can be simulated also with local external fields. However, in unmodulated (i.e. non engineered) systems, which are easier to create experimentally, such a difference is fundamental. Motivated by this, in this paper we focus on unmodulated qubit networks

described by the XYZ Hamiltonian and we find a full characterization of the chains admitting pretty good state transfer, namely we prove the following result:

Theorem 1. *Pretty good state transfer occurs between the extremal vertices of Heisenberg chain of n qubits if and only if n is a power of 2. Moreover, in these cases, pretty good state transfer occurs between vertices at the j th and $(n + 1 - j)$ th position for all $j = 1, \dots, n$.*

The remainder of the paper is organized as follows. In Section II we introduce the required notation. Section III describes the tools needed for the proof of our main result. Specifically Theorem 2 generalizes for every symmetric algebraic matrix a result by Vinet and Zhedanov⁸ on certain XY spin chains. Finally Section IV proves our main result.

II. NOTATION AND DEFINITIONS

We consider a graph $G = (V, E)$ with a set of vertices $V(G) = \{1, \dots, n\}$ and a set of edges $E(G)$ that describe the physical pairwise couplings between two vertices. We denote $A(G)$ the adjacency matrix with elements $[A(G)]_{ij} = 1$, if $(i, j) \in E(G)$, and $[A(G)]_{ij} = 0$ otherwise. For a generic graph structure the Heisenberg (XYZ) Hamiltonian is defined by

$$\mathcal{H}_{\text{XYZ}}(G) = \frac{1}{2} \sum_{i \neq j} A(G)_{ij} (X_i X_j + Y_i Y_j + Z_i Z_j), \quad (1)$$

where X_i, Y_i, Z_i are the Pauli matrices acting on the i -th vertex. On the other hand, the XY Hamiltonian is

$$\mathcal{H}_{\text{XY}}(G) = \frac{1}{2} \sum_{i \neq j} A(G)_{ij} (X_i X_j + Y_i Y_j). \quad (2)$$

Both $\mathcal{H}_{\text{XY}}(G)$ and $\mathcal{H}_{\text{XYZ}}(G)$ act on the Hilbert space $(\mathbb{C}^2)^{\otimes n}$. We call $\{|0\rangle, |1\rangle\}$ the basis of the Pauli matrices on each vertex and we define the single-particle subspace as the Hilbert space generated by the vectors $X_i |0\rangle^{\otimes n} = |0 \dots 0 1 0 \dots 0\rangle \in (\mathbb{C}^2)^{\otimes n}$, for $i = 1, \dots, n$, where the $|1\rangle$ state is in the i -th position. Within this single-particle subspace, the above Hamiltonians can be written as (see Ref.⁶)

$$\mathcal{H}_{\text{XYZ}}^{(1)}(G) = |E(G)| \mathbb{1} - 2L(G), \quad (3)$$

$$\mathcal{H}_{\text{XY}}^{(1)}(G) = 2A(G), \quad (4)$$

where the subscript (1) refers to the single-particle subspace, $L(G) = \Delta(G) - A(G)$ is the Laplacian of the graph and $\Delta(G)$ is the diagonal matrix whose diagonal i -th entry is the

degree $d(i)$ of vertex i , namely the number of edges incident with i . For simplicity, we avoid the use of the explicit notation $\mathcal{H}_{\text{XYZ}}^{(1)}(G)$, $\mathcal{H}_{\text{XY}}^{(1)}(G)$, and we simply call $A(G)$ and $L(G)$ as the XY and XYZ Hamiltonians, as they are equivalent to equations ((1)) and ((2)) in the single-particle subspace up to a trivial rescaling and shift.

We now introduce the concept of perfect and pretty good state transfer. Given M a symmetric matrix whose columns are indexed by the set of vertices V , we say that *perfect state transfer* occurs between vertices a and b of M if there is a $\tau \in \mathbb{R}^+$ such that

$$|\exp(i\tau M)_{a,b}| = 1.$$

This framework generalizes the concept of state transfer in the quantum walk of XY and XYZ Hamiltonians in the single-excitation subspace where M is respectively chosen as $A(G)$ or $L(G)$. If it is clear from the context which M we are dealing with, we use the notation $\exp(itM) = U(t)$.

We relax the definition of perfect state transfer to an ϵ -version. We say that M admits *pretty good state transfer* (also known as almost perfect state transfer) between vertices a and b if, for any $\epsilon > 0$, there is a time $\tau > 0$ such that

$$|U(\tau)_{a,b}| > 1 - \epsilon, \tag{5}$$

If $\mathbf{e}_a$ and $\mathbf{e}_b$ are the characteristic vectors of columns a and b , equation (5) is equivalent to the existence of a $\lambda \in \mathbb{C}$ of absolute value equal to 1 such that

$$\|U(\tau)\mathbf{e}_a - \lambda\mathbf{e}_b\| < \epsilon.$$

Finally, for shortness, when ϵ is not relevant, we abbreviate this equation to

$$U(\tau)\mathbf{e}_a \approx \lambda\mathbf{e}_b.$$

Given a real symmetric matrix M with spectral decomposition

$$M = \sum_{r=0}^d \theta_r E_r,$$

we say that a and b are *strongly cospectral* if $E_r\mathbf{e}_a = \pm E_r\mathbf{e}_b$ for all r . This nomenclature is inspired by the following fact. We say that vertices a and b are *cospectral* if the matrix obtained from M upon removing row and column indexed by a has the same spectrum as

when we remove row and column indexed by b . An equivalent formulation is that $(E_r)_{a,a} = (E_r)_{b,b}$ for all r , therefore every pair of strongly cospectral vertices is cospectral, as one would expect. If M is either the adjacency or the Laplacian matrix of a graph, cospectral vertices have necessarily the same number of neighbours. Moreover, in the adjacency case, a and b are cospectral if and only if, for all $k \in \mathbb{Z}$, the number of walks of length k that start and end in a is the same as the number for b (see Ref.²² (Section 2.5) for proofs and references of these facts). There are cases in which cospectral vertices are not strongly cospectral, and in fact we do not know any combinatorial characterization of this property. Finally, it is worth mentioning that if all eigenvalues are simple, both properties are equivalent, and that if M is a tridiagonal matrix (thus encoding the adjacency of a linear chain), then strong cospectrality is equivalent to the property of mirror-symmetry of the weights.

We also define the *eigenvalue support* of a as the set of eigenvalues θ_r such that $E_r \mathbf{e}_a \neq 0$.

III. TECHNICAL PRELIMINARIES

Godsil et al.⁹ determined when a linear chain with unmodulated spins admits pretty good state transfer between the end vertices according to the XY-Hamiltonian. Subsequently, Vinet and Zhedanov⁸ applied a theorem due to Kronecker on chains with non-unitary weights, providing new examples of pretty good state transfer in the XY-Hamiltonian model.

We point out that the application of Kronecker's theory to characterize pretty good state transfer in Heisenberg chains of prime length was previously considered¹⁴. Unfortunately the fact that the eigenvalues of these Hamiltonians are not linearly independent over the rationals went unnoticed, leading to the false conclusion that pretty good state transfer would occur in all chains of prime length. As an example, note that the probability of state transfer at time t between the extreme vertices in a chain of 3 qubits is equal to $\frac{1}{9}(1 - \cos(t))^2(5 + 4\cos(t))$, which is at most $3/4$ for all t .

In this work, we apply a more descriptive version of Kronecker's theorem, which deals with the case where the eigenvalues are not necessarily linearly independent over the rationals. As a result, we fully characterize linear chains with unmodulated spins admitting pretty good state transfer according to the Heisenberg Hamiltonian.

Theorem 2. *Let a and b be columns of a symmetric algebraic matrix M . Then pretty good state transfer occurs between a and b if and only if both conditions below are satisfied.*

- (i) *Columns a and b are strongly cospectral. In this case, let $\theta_0, \dots, \theta_d$ be the eigenvalues in their support, and for $r = 0, \dots, d$, let σ_r be defined as 0 if the projections onto E_r are equal, and 1 if they have opposite signs.*
- (ii) *If there is a set of integers $\ell_0, \dots, \ell_d$ such that*

$$\sum_{r=0}^d \ell_r \theta_r = 0 \quad \text{and} \quad \sum_{r=0}^d \ell_r \sigma_r \text{ is odd,}$$

then

$$\sum_{r=0}^d \ell_r \neq 0.$$

Condition (i) is known to be necessary for perfect state transfer (see for instance Ref.²³). In the Lemma below, a slightly modified argument works to show that it is also necessary for pretty good state transfer.

Lemma 3. *If pretty good state transfer occurs between a and b , then they are strongly cospectral vertices.*

Proof. From the spectral decomposition, we have

$$U(t) = \sum_{r=0}^d e^{it\theta_r} E_r,$$

thus

$$|U(t)_{a,b}| \leq \sum_{r=0}^d |(E_r)_{a,b}|.$$

Now $\sum E_r = I$, and, by Cauchy-Schwartz,

$$(E_r)_{a,a} \geq |(E_r)_{a,b}|.$$

Thus

$$\sum_{r=0}^d |(E_r)_{a,b}| \leq 1$$

and equality holds if and only if, for all r ,

$$(E_r)_{a,a} = |(E_r)_{a,b}|,$$

or equivalently, a and b are strongly cospectral. Therefore if a and b are not strongly cospectral, then there is ϵ such that

$$\sum_{r=0}^d |(E_r)_{a,b}| < 1 - \epsilon,$$

and so, for all t , it follows that

$$|U(t)_{a,b}| \leq 1 - \epsilon,$$

hence pretty good state transfer does not occur. $\square$

We will make use of the following result due to Kronecker.

Theorem 4 (Kronecker, see for instance Ref.²⁴, Chapter 3). *Let $\theta_0, \dots, \theta_d$ and $\zeta_0, \dots, \zeta_d$ be arbitrary real numbers. For an arbitrarily small ϵ , the system of inequalities*

$$|\theta_r y - \zeta_r| < \epsilon \pmod{2\pi}, \quad (r = 0, \dots, d),$$

admits a solution for y if and only if, for integers $\ell_0, \dots, \ell_d$, if

$$\ell_0 \theta_0 + \dots + \ell_d \theta_d = 0,$$

then

$$\ell_0 \zeta_0 + \dots + \ell_d \zeta_d \equiv 0 \pmod{2\pi}.$$

Now we prove our characterization.

Proof of Theorem 2. Observe that

$$U(\tau) \mathbf{e}_a \approx \lambda \mathbf{e}_b$$

is equivalent to, for all r ,

$$e^{i\theta_r \tau} E_r \mathbf{e}_a \approx \lambda E_r \mathbf{e}_b,$$

which in turn, when $\lambda = e^{i\delta}$, is equivalent to, for all r such that $E_r \mathbf{e}_u \neq 0$,

$$\theta_r \tau \approx \delta + q_r \pi, \tag{6}$$

where $q_r \in \mathbb{Z}$ is even if and only if $E_r \mathbf{e}_u = E_r \mathbf{e}_v$, and odd if and only if $E_r \mathbf{e}_u = -E_r \mathbf{e}_v$.

A solution to equation (6) is equivalent to a solution as described in Theorem 4 with

$$y = \tau \quad \text{and} \quad \zeta_r = \delta + \sigma_r \pi,$$

where $\sigma_r = 0$ if $E_r \mathbf{e}_a = E_r \mathbf{e}_b$, and $\sigma_r = 1$ if $E_r \mathbf{e}_a = -E_r \mathbf{e}_b$.

Now, a particular set of integers $\ell_0, \dots, \ell_d$ satisfies

$$\ell_0 \zeta_0 + \dots + \ell_d \zeta_d \equiv 0 \pmod{2\pi}$$

if and only if there is a δ such that

$$\ell_0(\delta + \sigma_0\pi) + \dots + \ell_d(\delta + \sigma_d\pi) \equiv 0 \pmod{2\pi}$$

which in turn is equivalent to

$$\delta \left(\sum_{r=0}^d \ell_r \right) + \pi \left(\sum_{r=0}^d \sigma_r \ell_r \right) \equiv 0 \pmod{2\pi}. \quad (7)$$

A solution δ to this equation exists if and only if, if $\sum \sigma_r \ell_r$ is odd, then $\sum \ell_r$ is non-zero, precisely as stated in condition (ii). This shows immediately that condition (ii) is necessary.

Note however that in order to apply Kronecker's Theorem and show sufficiency, we still need to show that the δ chosen to solve (7) should also work for any other set of integers $\ell'_0, \dots, \ell'_d$ satisfying $\sum \ell'_r \theta_r = 0$. The result follows easily if, for all such sets, $\sum \sigma_r \ell'_r$ is even, in which case $\delta = 0$ is a solution. So assume $\ell_0, \dots, \ell_d$ are such that $\sum \sigma_r \ell_r$ is odd, and so $\delta \sum \ell_r$ is also odd, and let $\ell'_0, \dots, \ell'_d$ be such that $\sum \ell'_r \theta_r = 0$. Let $\alpha = \sum \ell_r$ and $\beta = \sum \ell'_r$, and define integers $\gamma_r = (\delta/\pi)(\beta \ell_r - \alpha \ell'_r)$. If $\sum \sigma_r \ell'_r$ and $(\delta/\pi) \sum \ell'_r$ have opposing parities, then $\sum \gamma_r \theta_r = 0$, $\sum \sigma_r \gamma_r$ is odd, and $\sum \gamma_r = 0$. This contradicts condition (ii). Therefore $\sum \sigma_r \ell'_r$ and $(\delta/\pi) \sum \ell'_r$ have the same parity, and thus δ and $\ell'_0, \dots, \ell'_d$ satisfy (7) as well. $\square$

This next corollary is notably useful to study the Laplacian matrix.

Corollary 5. *Assume that 0 is an eigenvalue of M in the support of strongly cospectral columns a and b . Say the other eigenvalues in their support are $\theta_1, \dots, \theta_d$, and have $\sigma_1, \dots, \sigma_d$ defined as before. Then pretty good state transfer occurs between a and b if and only if whenever there are integers $\ell_1, \dots, \ell_d$ such that*

$$\sum_{r=1}^d \ell_r \theta_r = 0,$$

then

$$\sum_{r=1}^d \sigma_r \ell_r \text{ is even.}$$

Moreover, in this case, the complex phase with which pretty good state transfer occurs will be equal to 1.

Proof. Make $\theta_0 = 0$. Then given $\ell_0, \dots, \ell_d$,

$$\sum_{r=0}^d \ell_r \theta_r = 0 \quad \Longleftrightarrow \quad \sum_{r=1}^d \ell_r \theta_r = 0.$$

Hence the choice of ℓ_0 is arbitrary, and thus can always be made such that

$$\sum_{r=0}^d \ell_r = 0.$$

Thus, in order for pretty good state transfer to occur, $\sum_{r=1}^d \sigma_r \ell_r$ can never be odd, and if it is even in all cases, condition (ii) of Theorem 2 is vacuously satisfied. Moreover, in this case, as the choice ℓ_0 is arbitrary and hence can also be made in a way that $\sum_{r=0}^d \ell_r$ is odd, δ must be an even multiple of π , therefore $\lambda = e^{i\delta} = 1$. $\square$

A. The spectrum of Heisenberg chains

We refer the reader to Brouwer and Haemers²⁵ for the result below. Let P_n denote the path on n vertices. Recall that $L(X)$ denotes the Laplacian matrix of the graph X .

- The eigenvalues of $L(P_n)$ are 0 with the all 1s eigenvector, and $2 + 2\cos(\pi r/n)$, $r = 1, \dots, n-1$. If $\beta_k = \sin(k\pi r/n)$, its corresponding eigenvector is

$$(\beta_1, (-1)^1(\beta_1 + \beta_2), (\beta_2 + \beta_3), \dots, (-1)^n(\beta_{n-2} + \beta_{n-1}), (-1)^{n+1}\beta_{n-1}).$$

IV. MAIN RESULT

We are ready to prove Theorem 1.

Theorem 1 (restated) *Pretty good state transfer occurs on $L(P_n)$ between the extremal vertices if and only if n is a power of 2. Moreover, in these cases, pretty good state transfer occurs between vertices at the j th and $(n+1-j)$ th position for all $j = 1, \dots, n$.*

Proof. Suppose the spectral decomposition of $L(P_n)$ is given by

$$L(P_n) = \sum_{r=0}^{n-1} \lambda_r E_r.$$

Let R be the anti-diagonal matrix of order n . It is a straightforward consequence of the spectrum of P_n described in Section III A that

$$\sum_{r=0}^{n-1} (-1)^r E_r = R.$$

This readily implies that vertices at positions j and $(n+1-j)$ are strongly cospectral for $j = 1, \dots, n$, and hence condition (i) of Theorem 2 is always satisfied, with $\sigma_r = (-1)^r$.

Let $\zeta_{2n} = e^{\pi/n}$. Clearly the eigenvalues of P_n can be expressed as

$$\lambda_r = 2 - (\zeta_{2n}^r + \overline{\zeta_{2n}^r}) = 2 - (\zeta_{2n}^r + \zeta_{2n}^{2n-r}).$$

As a consequence, the eigenvalues belong to the cyclotomic field of ζ_{2n} . Now assume there are integers $\ell_1, \dots, \ell_{n-1}$ such that

$$\sum_{r=1}^{n-1} \ell_r (-2 + (\zeta_{2n}^r + \zeta_{2n}^{2n-r})) = 0. \quad (8)$$

If $\ell_0 = -\sum_{r=1}^{n-1} \ell_r$, then the cyclotomic polynomial $\Phi_{2n}(x)$ divides

$$L(x) = 2\ell_0 + \sum_{r=1}^{n-1} \ell_r x^r + \sum_{r=n+1}^{2n-1} \ell_{2n-r} x^r. \quad (9)$$

- If n is a power of 2, then $\Phi_{2n}(x) = 1 + x^n$. Performing long division starting from the terms of smaller degree, the general form of an exact quotient of the division of $L(x)$ by $\Phi_{2n}(x)$ is

$$2\ell_0 + \sum_{r=1}^{n-1} \ell_r x^r,$$

thus the division is exact (and equation (8) is satisfied) if and only if $\ell_0 = 0$ and $\ell_s = \ell_{n-s}$ for all $s = 1, \dots, n$. As a consequence, whenever (8) holds, $\sum \ell_{\text{odd}}$ is always even, and pretty good state transfer occurs.

- If n is an odd prime, then $\Phi_{2n}(x) = 1 - x + x^2 - \dots + x^{n-1}$. Performing long division starting from the terms of smaller degree, the general form of an exact quotient of the division of $L(x)$ by $\Phi_{2n}(x)$ is

$$2\ell_0 + (2\ell_0 + \ell_1)x + \sum_{r=2}^{n-1} (\ell_r + \ell_{r-1})x^r + \ell_1 x^n.$$

This implies that a set of integers $\ell_1, \dots, \ell_{n-1}$ satisfy equation (8) if and only if, for all odd s between 1 and $n-1$,

$$\ell_s - \ell_{n-s} = -2\ell_0 = 2 \sum_{r=1}^{n-1} \ell_r. \quad (10)$$

If $n \equiv 3 \pmod{4}$, then, for any odd s , define $\ell_s = n$ and $\ell_{n-s} = -(n-2)$. This provides a solution to (10) such that

$$\sum_{s \text{ odd}} \ell_s \text{ is odd,}$$

hence pretty good state transfer does not occur in this case.

If $n \equiv 1 \pmod{4}$, make $\ell_1 = -6$, $\ell_{n-1} = 2(n-2)$, and, for any other odd s , $\ell_s = -(n+4)$ and $\ell_{n-s} = (n-2)$. Hence

$$\sum_{r=1}^{n-1} \ell_r = -(n+1),$$

and so, for all s odd,

$$\ell_s - \ell_{n-s} = 2 \sum_{r=1}^{n-1} \ell_r,$$

but

$$\sum_{s \text{ odd}} \ell_s = -(n+4) \left(\frac{n-3}{2} \right) - 6 \text{ is odd,}$$

hence pretty good state transfer does not occur in this case.

- Now suppose $n = mj$, with m odd, $m > 1$ and $j > 1$. Let p be an odd prime dividing m , thus $2n = pk$, with k a positive even number. It follows that $x^{2n} - 1 = \Phi_p(x)R(x)$ where

$$R(x) = \sum_{t=0}^{k-1} -x^{tp} + x^{tp+1}.$$

Let ℓ_r be defined as 0, 1 or -1 in such way that

$$R(x) = \sum_{r=0}^{n-1} \ell_r x^r.$$

Note that 1, ζ_{2n} and ζ_{2n}^{2n-1} are all roots of $R(x)$. Thus

$$R(\zeta_{2n}) + R(\zeta_{2n}^{2n-1}) = 0 \quad \text{and} \quad \sum_{r=1}^{n-1} \ell_r = 1.$$

Hence

$$0 = R(\zeta_{2n}) + R(\zeta_{2n}^{2n-1}) = \sum_{r=1}^{n-1} \ell_r (-2 + \zeta_{2n}^r + \zeta_{2n}^{2n-r})$$

where, for $t = 1, \dots, k-1$,

$$\ell_r = 1 \text{ if } r = tp + 1, \quad \ell_r = -1 \text{ if } r = tp, \quad \ell_r = 0 \text{ otherwise.}$$

Note that t is odd if and only if tp is odd, and t is even if and only if $tp + 1$ is odd. Thus the sum of the ℓ_r with odd r is equal to -1 . Therefore pretty good state transfer does not occur in this case.

□

V. FINAL REMARKS

Theorem 2 provides conditions which are necessary and sufficient for pretty good state transfer to occur, but fails to offer an estimate of how large t should be in order to obtain an approximation in terms of a given ϵ . In Ref.²⁶, it is shown that in certain dynamical systems the time is proportional to $(1/\epsilon)^n$, but this makes the assumption that the numbers $\theta_0, \dots, \theta_d$ in the statement of Kronecker's Theorem are linearly independent over the rationals. Note that this is not the case for the examples studied in our paper, and we do not know a reference for these situations.

We point out that we are not necessarily determining all cases in which a qubit chain might admit pretty good state transfer, as we are focusing only on transfer between the end vertices. In fact, there are known examples of perfect state transfer in linear chains with weighted edges between internal nodes while not occurring between the end vertices (see Ref.⁴).

In an earlier version of our paper, we left as an open question to characterize pretty good state transfer between any pair of vertices in a linear chain with unit weights according to the XY or the XYZ Hamiltonian. For the XY Hamiltonian, some progress was made in Ref.²⁷, and the problem was subsequently solved in Ref.²⁸. The question remains open for the Laplacian case.

ACKNOWLEDGMENTS

LB acknowledges funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement No. 308253. GC acknowledges the support of grants FAPESP 15/16339-2 and FAPESP 13/03447-6. CG acknowledges the support of NSERC Grant RGPIN-9439. SS acknowledges the Royal Society and EPSRC.

We all acknowledge the valuable comments of the anonymous referee. We also acknowledge Gabor Lippner, who offered valuable insights about Theorem 2 that lead us to substantially improve its statement.

REFERENCES

- ¹Thaddeus D Ladd, Fedor Jelezko, Raymond Laflamme, Yasunobu Nakamura, Christopher Monroe, and Jeremy L O'Brien. Quantum computers. *Nature*, 464(7285):45–53, 2010.
- ²Sougato Bose. Quantum Communication through an Unmodulated Spin Chain. *Physical Review Letters*, 91(20):207901, nov 2003.
- ³Georgios M Nikolopoulos and Igor Jex (Eds.). *Quantum State Transfer and Network Engineering*. Springer, 2014.
- ⁴Alastair Kay. Perfect, efficient, state transfer and its application as a constructive tool. *International Journal of Quantum Information*, 8(04):641–676, 2010.
- ⁵Chris D Godsil. When can perfect state transfer occur? *Electronic Journal of Linear Algebra*, 23:877–890, 2012.
- ⁶Sougato Bose, Andrea Casaccino, Stefano Mancini, and Simone Severini. Communication in XYZ all-to-all quantum networks with a missing link. *International Journal of Quantum Information*, 7(04):713–723, 2009.
- ⁷Daniel Burgarth and Sougato Bose. Conclusive and arbitrarily perfect quantum-state transfer using parallel spin-chain channels. *Physical Review A*, 71(5):052315, 2005.
- ⁸Luc Vinet and Alexei Zhedanov. Almost perfect state transfer in quantum spin chains. *Physical Review A*, 86(5):052319, nov 2012.
- ⁹Chris D Godsil, Stephen Kirkland, Simone Severini, and Jamie Smith. Number-theoretic nature of communication in quantum spin systems. *Physical Review Letters*, 109(5):050502,

aug 2012.

- ¹⁰Leonardo Banchi. Ballistic quantum state transfer in spin chains: General theory for quasi-free models and arbitrary initial states. *The European Physical Journal Plus*, 128(11):1–18, 2013.
- ¹¹Tony JG Apollaro, Leonardo Banchi, Alessandro Cuccoli, Ruggero Vaia, and Paola Verrucchi. 99%-fidelity ballistic quantum-state transfer through long uniform channels. *Physical Review A*, 85(5):052319, 2012.
- ¹²Salvatore Lorenzo, Tony JG Apollaro, Antonello Sindona, and Francesco Plastina. Quantum-state transfer via resonant tunneling through local-field-induced barriers. *Physical Review A*, 87(4):042313, 2013.
- ¹³Yasser Omar and Rúben Sousa. Pretty good state transfer of entangled states through quantum spin chains. *New Journal of Physics*, 16(12):123003, 2014.
- ¹⁴Daniel Burgarth. *Quantum State Transfer with Spin Chains*. PhD thesis, University of London, 2007.
- ¹⁵L. Campos Venuti, C. Degli Esposti Boschi, and M. Roncaglia. Qubit Teleportation and Transfer across Antiferromagnetic Spin Chains. *Physical Review Letters*, 99:060401, 2007.
- ¹⁶L. Campos Venuti, S. M. Giampaolo, F. Illuminati, and P. Zanardi. Long-distance entanglement and quantum teleportation in XX spin chains. *Physical Review A*, 76:052328, 2007.
- ¹⁷Hans-Jürgen Mikeska, Alexei K. Kolezhuk. One-dimensional magnetism, Quantum magnetism. Springer Berlin Heidelberg, 2004. 1-83.
- ¹⁸Claudio Albanese, Matthias Christandl, Nilanjana Datta, and Artur Ekert. Mirror inversion of quantum states in linear registers. *Physical Review Letters*, 93(23):230502, 2004.
- ¹⁹Ronald Hanson, Leo P Kouwenhoven, Jason R Petta, Seigo Tarucha, and Lieven MK Vandersypen. Spins in few-electron quantum dots. *Reviews of Modern Physics*, 79(4):1217, 2007.
- ²⁰Bruce E Kane. A silicon-based nuclear spin quantum computer. *Nature*, 393(6681):133–137, 1998.
- ²¹Takeshi Fukuhara, Adrian Kantian, Manuel Endres, Marc Cheneau, Peter Schauß, Sebastian Hild, David Bellem, Ulrich Schollwöck, Thierry Giamarchi, Christian Gross, et al. Quantum dynamics of a mobile spin impurity. *Nature Physics*, 9(4):235–241, 2013.
- ²²Gabriel Coutinho. *Quantum State Transfer in Graphs*. PhD dissertation, University of

Waterloo, 2014.

- ²³Kay, A. Basics of perfect communication through quantum networks. *Physical Review A*, 84(2), 022337, 2011.
- ²⁴Boris M Levitan and Vasilii V Zhikov. *Almost Periodic Functions and Differential Equations*. CUP Archive, 1982.
- ²⁵Andries E Brouwer and Willem H Haemers. *Spectra of Graphs*. Universitext. Springer, New York, 2012.
- ²⁶P Chr Hemmer, L C Maximon, and H Wergeland. Recurrence Time of a Dynamical System *Physical Review*, 111:689–694, 1957.
- ²⁷Gabriel Coutinho, Krystal Guo, and Christopher van Bommel. Pretty good state transfer between internal nodes of paths. *Arxiv*, 1611.09836.
- ²⁸Christopher van Bommel. A Complete Characterization of Pretty Good State Transfer on Paths. *Arxiv*, 1612.05603.